

PS2




My score

90.6% (14.5/16)

Q1


4 / 4

1. Two springs each have spring constant k and equilibrium length l . They are both stretched a distance l and attached to a mass m and two walls, as shown in figure below. At a given instant, the right spring constant is somehow magically changed to $3k$ (the relaxed length remains l). What is the resulting $x(t)$? Take the initial position to be $x = 0$.



PH254 PS2

Q1/



$$\vec{F}_1 = -K(l+x) \quad [\text{LEFT}]$$

$$\vec{F}_2 = -3K(l-x) \quad [\text{RIGHT}]$$

$$\text{Then } F = ma = -K(l+x) + 3K(l-x)$$

$$\Rightarrow m\ddot{x} = -Kl - Kx + 3Kl - 3Kx \\ = 2Kl - 4Kx$$

• Find the homogeneous solution:

$$\ddot{x} + 4\frac{K}{m}x = 0$$

$$\text{Guess } y = Ae^{rt}$$

$$r^2 + 4\frac{K}{m} = 0$$

$$\Rightarrow r^2 = -4\frac{K}{m}$$

$$\Rightarrow r = \pm i2\sqrt{\frac{K}{m}}$$

$$\text{Thus our homogeneous solution is } x_h(t) = A_1 e^{i2\sqrt{\frac{K}{m}}t} + A_2 e^{-i2\sqrt{\frac{K}{m}}t}$$

• Find a particular solution:

$$\ddot{x} + 4\frac{K}{m}x = 2\frac{K}{m}l \quad \text{If } x_p(t) \text{ is a constant, } \ddot{x}_p = 0.$$

$$\Rightarrow \frac{4}{m}x = 2\frac{l}{m} \Rightarrow x_p = \frac{1}{2}l$$

Our general solution is then

$$x = x_h + x_p = A_1 e^{i2\sqrt{\frac{K}{m}}t} + A_2 e^{-i2\sqrt{\frac{K}{m}}t} + \frac{1}{2}l$$



Q2

3 / 4

2. (a) Two vibrations are given by:

$$x_1 = A \cos(\omega_1 t + \phi_1) \text{ and } x_2 = A \cos(\omega_2 t + \phi_2).$$

Find an expression for their superposed motion in the form

$$a \cos(bt + c) \cos(dt + e)$$

and determine the constants a through e in terms of the given parameters. Use complex notation in your derivation, as in the class notes.

- (b) Consider the vibrations $x_1 = 3 \cos(3t + \pi/5)$ and $x_2 = 4 \cos(4t + \pi/8)$.

What is the expected period for $x_1 + x_2$?

- (c) **Complex number and phasor practice:** Express the following in the form $z = \operatorname{Re}[Ae^{i(\omega t + \alpha)}]$ and sketch them by hand or plot them in pylab in the complex plane at $t = \pi/\omega$ (i.e. at one half period):

$$z = 5 \sin \omega t + 6 \cos \omega t;$$

$$z = \sin(\omega t + \pi/3) - \cos \omega t;$$

$$z = \sin \omega t + 2 \cos(\omega t + \pi/6) + \cos \omega t.$$

note: just sketch by hand is fine

For the sketch, you should sketch

$Ae^{i(\omega t + \alpha)}$ (not z , since z is real) in the complex plane at $t = \pi/\omega$

PHY254 P52

Q2) 2) $x_1 = A \cos(\omega_1 t + \phi_1)$, $x_2 = A \cos(\omega_2 t + \phi_2)$.

→ Same Amplitude!

$$x_1 = \operatorname{Re} \{ A e^{i(\omega_1 t + \phi_1)} \}, \quad x_2 = \operatorname{Re} \{ A e^{i(\omega_2 t + \phi_2)} \}.$$

$$\text{Then } x_1 + x_2 = \operatorname{Re} \{ A e^{i(\omega_1 t + \phi_1)} + A e^{i(\omega_2 t + \phi_2)} \}.$$

$$\text{Let } \omega_1 t + \phi_1 = \alpha + \beta, \quad \omega_2 t + \phi_2 = \alpha - \beta.$$

$$\Rightarrow \alpha = \omega_1 t + \phi_1 - \beta$$

$$\omega_2 t + \phi_2 = \omega_1 t + \phi_1 - 2\beta \Rightarrow \beta = -\frac{1}{2} [(\omega_2 - \omega_1)t + (\phi_2 - \phi_1)]$$

$$\Rightarrow \alpha = \frac{1}{2} [(\omega_2 + \omega_1)t + (\phi_2 + \phi_1)].$$

$$\text{Let } \omega_2 - \omega_1 = \Delta\omega, \quad \phi_2 - \phi_1 = \Delta\phi.$$

$$\text{And } \omega_2 + \omega_1 = \delta\omega, \quad \phi_2 + \phi_1 = \delta\phi.$$

$$\Rightarrow x_1 + x_2 = \operatorname{Re} \{ A e^{i(\alpha + \beta)} + A e^{i(\alpha - \beta)} \} = \operatorname{Re} \{ A e^{i\alpha} (e^{i\beta} + e^{-i\beta}) \}$$

$$\text{However } e^{i\beta} + e^{-i\beta} = \cos(\beta) + i\sin(\beta) + \cos(\beta) - i\sin(\beta) \\ = 2\cos(\beta).$$

$$= \operatorname{Re} \left\{ A e^{\frac{i}{2}\delta\omega t} e^{\frac{i}{2}\delta\phi} \left[2 \cos\left(-\frac{1}{2}\Delta\omega t - \frac{1}{2}\Delta\phi\right) \right] \right\}$$

$$= \operatorname{Re} \left\{ A \left(\cos\left(\frac{\delta\omega}{2}t\right) + i\sin\left(\frac{\delta\omega}{2}t\right) \right) \left(\cos\left(\frac{\delta\phi}{2}\right) + i\sin\left(\frac{\delta\phi}{2}\right) \right) \dots \right\}$$

$$= \operatorname{Re} \left\{ 2A \underbrace{\left(\cos\left(\frac{\delta\omega}{2}t\right)\cos\left(\frac{\delta\phi}{2}\right) - \sin\left(\frac{\delta\omega}{2}t\right)\sin\left(\frac{\delta\phi}{2}\right) \right)}_{= \cos\left(\frac{\delta\omega}{2}t + \frac{\delta\phi}{2}\right)} \right\} \quad \text{[trig/havy ... terms]}$$

$$= 2A \cos\left(\frac{\delta\omega}{2}t + \frac{\delta\phi}{2}\right) \cos\left(-\frac{\Delta\omega}{2}t + -\frac{\Delta\phi}{2}\right)$$

Thus

$$x_1 + x_2 = 2A \cos\left(\frac{\omega_1 + \omega_2}{2}t + \frac{\phi_1 + \phi_2}{2}\right) \cos\left(\frac{\omega_1 - \omega_2}{2}t + \frac{\phi_1 - \phi_2}{2}\right)$$

b) Consider $x_1 = 3\cos(3t + \frac{\pi}{5})$, $x_2 = 4\cos(4t + \frac{\pi}{8})$

Expected period $T_{ex} = \frac{2\pi}{\omega_{ex}}$.

Notice that $\omega_1 = 3$, $\omega_2 = 4$

$\Rightarrow \frac{\omega_1}{\omega_2} = \frac{3}{4} = \frac{m}{n} = r$,

so ω_1 & ω_2 are commensurate.

This implies that $x_1 + x_2$ has a periodic superposition.

$mT_1 = 3 \cdot \frac{2\pi}{3} = 2\pi$, and

$nT_2 = 4 \cdot \frac{2\pi}{4} = 2\pi$,

Therefore $T_{ex} = 2\pi$.

c) Find and plot the phasor diagrams in the form $z = \text{Re}\{Ae^{i(\omega t + \phi)}\}$.


i) $z_1 = 5\sin \omega t + 6\cos \omega t$.

$= \text{Re}\{5(-i)e^{i(\omega t)} + 6e^{i(\omega t)}\} = \text{Re}\{(6-5i)e^{i\omega t}\}$.

Express $6-5i$ as some $e^{i\phi}$. $r = \sqrt{6^2 + (-5)^2} = \sqrt{61}$,


and $\phi = \arctan(-\frac{5}{6}) = -\arctan(\frac{5}{6})$; thus

$z_1 = \text{Re}\{\sqrt{61}e^{i(\omega t - \arctan(5/6))}\}$



This is not
at
 $t = \pi/\omega$

$$\begin{aligned}
 \text{ii) } z_2 &= \sin(\omega t + \frac{\pi}{3}) - \cos \omega t \\
 &= \operatorname{Re} \{ -i e^{i(\omega t + \pi/3)} \} - \operatorname{Re} \{ e^{i\omega t} \} \\
 &= \operatorname{Re} \{ (-1-i) e^{i\omega t} e^{i\pi/3} \} \\
 &= \cos \frac{\pi}{3} + i \sin \frac{\pi}{3} = \frac{1}{2} + i \frac{\sqrt{3}}{2} \\
 &= \operatorname{Re} \{ (-1-i) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) e^{i\omega t} \} \\
 &= \operatorname{Re} \left\{ \left(\frac{\sqrt{3}-1}{2} - i \frac{\sqrt{3}+1}{2} \right) e^{i\omega t} \right\} \\
 r &= \sqrt{\left(\frac{\sqrt{3}-1}{2} \right)^2 + \left(\frac{\sqrt{3}+1}{2} \right)^2} = \sqrt{2} \\
 \varphi &= \arctan \left(- \frac{\sqrt{3}+1}{\sqrt{3}-1} \right)
 \end{aligned}$$




Thus $z_2 = \operatorname{Re} \left\{ \sqrt{2} e^{i(\omega t - \frac{\pi}{4})} \right\}$

Need to add pi to this

$$\begin{aligned}
 \text{iii) } z_3 &= \sin \omega t + 2 \cos(\omega t + \frac{\pi}{6}) + \cos \omega t \\
 &= \operatorname{Re} \{ -i e^{i\omega t} \} + \operatorname{Re} \{ 2 e^{i(\omega t + \pi/6)} \} + \operatorname{Re} \{ e^{i\omega t} \} \\
 &= \operatorname{Re} \{ (-i + 2 e^{i\pi/6} + 1) e^{i\omega t} \} \\
 &= \operatorname{Re} \{ (-i + 2(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6}) + 1) e^{i\omega t} \} = \operatorname{Re} \{ (\sqrt{3}+1) e^{i\omega t} \}
 \end{aligned}$$

Thus $z_3 = \operatorname{Re} \{ (\sqrt{3}+1) e^{i\omega t} \}$



Q3

4 / 4

3. A mass m connected to an ideal spring slides frictionlessly on a surface. It is connected to a “dashpot” damper that produces a resistive force proportional to the velocity v of the mass.




Figure 1: Damped oscillator

We did some experiments on this device and made the following observations:

- If the block is pushed horizontally to the left with a static force equal to mg , the spring compresses a distance h .
- If the block is moved at a steady speed $U = 3\sqrt{gh}$, the dashpot damper produces a resistive force of magnitude mg .

Now the system is displaced and released from rest, setting it into oscillation.

- Find the angular frequency of the underdamped oscillations of the system, expressed as a multiple of $\sqrt{g/h}$.
- After what time, expressed as a multiple of $\sqrt{h/g}$, is the amplitude of the underdamped motion decreased by a factor of e from its initial displacement?

PH4254 PS2

Q3) Forces: $F = -bv$ damping force
 $F = -kx$ restoring spring force.

$$F = mg = -k(-h), \Rightarrow k = \frac{mg}{h}. \quad \text{Similarly,}$$

$$F = mg = -b \cdot 3\sqrt{gh}, \Rightarrow b = \frac{m}{3} \sqrt{\frac{g}{h}}.$$

Our ODE is then $F = F_{sp} + F_{damp}$, so

$$m\ddot{x} = -\frac{mg}{h}x - \frac{m}{3}\sqrt{\frac{g}{h}}\dot{x}.$$

$$\text{This implies that } \ddot{x} + \frac{g}{h}x + \frac{1}{3}\sqrt{\frac{g}{h}}\dot{x} = 0.$$

a) Find angular frequency. We want a solution of the form

$x = e^{it} \cdot A$. Let $p = \sqrt{\frac{g}{h}}$ for simplicity. Guess $x_h = Ae^{rt}$. Then

$$r^2 + \frac{1}{3}pr + p^2 = 0. \Rightarrow 3r^2 + pr + 3p^2 = 0.$$

$$\text{Quadratic formula: } r = \frac{-p \pm \sqrt{p^2 - 4(3)(3p^2)}}{6} = \frac{-p \pm p\sqrt{1-36}}{6}$$

$$\Rightarrow r = -\frac{p}{6} \pm i \frac{p\sqrt{35}}{6}.$$

Our general solution is then

$$x = A e^{-\frac{p}{6}t} e^{i \frac{p\sqrt{35}}{6}t} + B e^{-\frac{p}{6}t} e^{-i \frac{p\sqrt{35}}{6}t}.$$

Since the angular frequency ω is given in the complex

solution $e^{i\omega t}$, this implies that $\omega = p \cdot \frac{\sqrt{35}}{6}$.

$$\text{Thus } \omega = \frac{\sqrt{35}}{6} \cdot \sqrt{\frac{g}{h}}$$

b) Time for amplitude to decay a e^{-1} factor:

In the solution to the ODE, the amplitude term is decayed by the term $e^{-\frac{b}{m}t} = e^{-\sqrt{\frac{g}{h}} \cdot \frac{t}{6}}$.

$\Rightarrow$ Solve for t in $e^{-\sqrt{\frac{g}{h}} \cdot \frac{t}{6}} = e^{-1}$,

which implies that

$$\sqrt{\frac{g}{h}} \cdot \frac{t}{6} = 1, \quad \text{or} \quad \boxed{t = 6 \cdot \sqrt{\frac{h}{g}}},$$

as desired.

Q4

3.5 / 4

4. Consider a particle of mass m , free to move in one dimension x , which is subjected to a drag force $-bv = -b\dot{x}$ but *no* spring-like restoring force. The particle is initially at rest, so $x_0 = x(0) = 0$ and $v_0 = \dot{x}(0) = 0$. At $t = 0$ we suddenly begin to apply a driving force $f(t) = mA \cos(\omega t)$ to the particle, where $A > 0$ is a constant.

For $t > 0$, the particle exhibits some transient motion but after a long time it settles down into a regular steady-state oscillation.

Note for (a): transient part means that we try to get the “homogeneous” linear differential equations, i.e. by assuming that the driving force is zero (the drag force still exists)

- (a) Write down $F = ma$ for this system and show that the *transient part* $x_t(t)$ of the total displacement $x(t)$ of the particle is given by

$$x_t(t) = C + \left(\frac{D}{\beta}\right)e^{-\beta t} \quad , \quad (1)$$

where $\beta = b/m$ and C and D are constants to be determined later from the initial conditions.

- (b) We learned in class that the long term, or steady state, part of the motion is found from the *particular solution* $x_p(t)$ of the forced equation of motion.

Find a particular solution by “complexifying” the equation of motion by assuming a complex applied force $\tilde{f}(t) = mAe^{i\omega t}$, so that $f = \Re[\tilde{f}]$, where $\Re[\cdot]$ means “take the real part”. Using a clever guess, find a complex expression $\tilde{x}_p(t)$ for the long-term, steady-state motion.

- (c) By taking the real part of the complex expression you found in the previous part, find the real steady state motion $x_p(t) = \Re[\tilde{x}_p(t)]$. Write it in the following form

$$x_p(t) = B \cos(\omega t + \phi) \quad , \quad (2)$$

finding expressions for the real amplitude B and phase ϕ .

HINT: It will be useful to write $\tilde{x}_p$ in the polar form $Be^{i\phi}e^{i\omega t}$ first. $Be^{i\phi}$ is the complex *phasor* of the motion. Draw a phasor diagram in the complex plane.

You may or may not recall that
 $\cos(\alpha + \delta) = \cos(\alpha)\cos(\delta) - \sin(\alpha)\sin(\delta)$.

- (d) Finally, combine the real transient and the steady-state motion to get the full motion for all time, $x(t) = x_t(t) + x_p(t)$. Apply the initial conditions and determine the constants C and D . You may assume the result of the previous part and just write them in terms of B and ϕ .

PHY254 PS2

Q4)

$$F_{\text{net}} = F_{\text{drive}} + F_{\text{drag}}$$

$$m\ddot{x} = mA \cos(\omega t) - b\dot{x}$$

$$\Rightarrow \ddot{x} + \frac{b}{m}\dot{x} = A \cos(\omega t)$$

a) Transient motion: no driving force

$$\ddot{x} + \frac{b}{m}\dot{x} = 0. \quad \text{Let } \beta = \frac{b}{m}.$$

Let $p = \dot{x}$, and guess $p = Ke^{rt}$ for a constant K .

$$\dot{p} + \beta p = 0 \Rightarrow Kr e^{rt} + \beta K e^{rt} = 0$$

$$\Rightarrow r + \beta = 0, \quad \text{thus } r = -\beta.$$

$$\text{Then } p = \dot{x} = K e^{-\beta t}.$$

$$x = \int \dot{x} dt = \int K e^{-\beta t} dt$$

$$= -\frac{K}{\beta} e^{-\beta t} + C. \quad \text{Let } D = -K, \text{ another constant. Then}$$

$$x_t(t) = C + \left(\frac{D}{\beta}\right) e^{-\beta t}, \quad \text{as required.}$$

b) Steady state motion: guess a solution

$$\ddot{x} + \beta \dot{x} = A \cos(\omega t) \rightsquigarrow \ddot{x} + \beta \dot{x} = A e^{i\omega t}$$

Complexified solution.

Guess $\tilde{x} = B e^{i(\omega t + \phi)}$, for some initial phase ϕ and constant B .

$$\text{Let } \alpha = B e^{i\phi}. \quad \text{Then } \tilde{x}(t) = \alpha e^{i\omega t}.$$

$$\Rightarrow \ddot{x} = \alpha \omega e^{i\omega t}$$

$$\text{and } \ddot{x} = -\alpha \omega^2 e^{i\omega t}$$

Our complex ODE then becomes

$$-\alpha \omega^2 e^{i\omega t} + i\beta \omega \alpha e^{i\omega t} = A e^{i\omega t}$$

$$\Rightarrow -\alpha \omega^2 + i\beta \omega \alpha = A$$


Now we can solve for α :

$$\alpha(i\beta\omega - \omega^2) = A \Rightarrow \alpha = \frac{A}{i\beta\omega - \omega^2} = B e^{i\phi}$$

Notice that $i\beta\omega - \omega^2$ is complex, hence it can be written in complex polar form $r e^{i\phi}$ for some r and ϕ .

Notice

$$r = \sqrt{A^2 + B^2} = \sqrt{\beta^2 \omega^2 + \omega^4}$$



$$\text{Thus } \phi = \arctan\left(-\frac{\beta\omega}{\omega^2}\right) = -\arctan\left(\frac{\beta}{\omega}\right).$$

This implies

$$\alpha = \frac{A}{r} e^{-i\phi} = \frac{A}{\omega \sqrt{\beta^2 + \omega^2}} e^{\arctan(\frac{\beta}{\omega})} = B e^{i\phi}$$

$$\text{Therefore } B = \frac{A}{\omega \sqrt{\beta^2 + \omega^2}} \text{ and } \phi = \arctan\left(\frac{\beta}{\omega}\right).$$

Therefore our solution is

$$\tilde{x}(t) = B e^{i(\omega t + \phi)}$$

where $B = \frac{A}{\omega \sqrt{\beta^2 + \omega^2}}$ and $\phi = \arctan\left(\frac{\beta}{\omega}\right)$

missing a minus sign

c) Our previous solution was

$$\tilde{x}_p(t) = B e^{i(\omega t + \varphi)}$$

where $B = \frac{A}{\omega \sqrt{p^2 + \omega^2}}$ and $\varphi = \arctan\left(\frac{p}{\omega}\right)$.

Expanding, we have


$$\tilde{x}_p(t) = B [\cos(\omega t + \varphi) + i \sin(\omega t + \varphi)]$$


so $x_p(t) = \operatorname{Re}\{\tilde{x}_p(t)\} = B \cos(\omega t + \varphi)$.

$$x_p(t) = B \cos(\omega t + \varphi)$$

where $B = \frac{A}{\omega \sqrt{p^2 + \omega^2}}$ and $\varphi = \arctan\left(\frac{p}{\omega}\right)$.

Phasor Diagram: Plot $B e^{i\varphi}$:





d) Constructing the solution:

$$x = x_p + x_t$$

$$= B \cos(\omega t + \varphi) + C + \frac{D}{\beta} e^{-\beta t}$$

$$x_0 = 0, \quad \dot{x}_0 = 0.$$

$$\begin{aligned} x \quad 0 &= B \overset{\cos \varphi}{\cos(\omega t + \varphi)} + C + \frac{D}{\beta} e^{-\beta t} \\ &= B \cos \varphi + C + \frac{D}{\beta} \end{aligned}$$

$$\begin{aligned} \dot{x} \quad 0 &= -B \omega \overset{\sin \varphi}{\sin(\omega t + \varphi)} - D e^{-\beta t} \\ &= -B \omega \sin \varphi - D \end{aligned}$$

$$\Rightarrow D = -B \omega \sin \varphi$$

$$\Rightarrow C = -B \cos \varphi + \frac{B \omega \sin \varphi}{\beta}$$

Thus our solution is

$$x(t) = B \cos(\omega t + \varphi) - B \cos \varphi + \frac{B \omega \sin \varphi}{\beta} - \frac{B \omega \sin \varphi}{\beta} e^{-\beta t}$$

$$\text{where } \beta = \frac{b}{m}, \quad B = \frac{A}{\omega \sqrt{\beta^2 + \omega^2}}, \quad \text{and } \varphi = \arctan\left(\frac{\beta}{\omega}\right)$$

